arXiv:1111.4629v1 [math.DS] 20 Nov 2011

HERMAN'S CONDITION AND SIEGEL DISKS OF POLYNOMIALS

ARNAUD CHÉRITAT AND PASCALE ROESCH

Abstract. Herman proved the presence of critical points on the boundary of Siegel disks of unicritical polynomials under some diophantine condition now called the Herman condition. We extend this result to polynomials with two critical points.

1. Introduction

Definition 1. *Let $\mathcal{H}$ be the set of real numbers $\theta \in \mathbb{R}$ such that every orientation preserving analytic circle diffeomorphism of rotation number θ is analytically conjugate to a rotation.*

Herman proved that $\mathcal{H}$ is non empty by showing that it contains all diophantine numbers [Her79]. Yoccoz characterized numbers in $\mathcal{H}$ in terms of their continued fraction expansion [Yoc02].

We call *Siegel disk* of a rational map f of degree at least 2 a maximal domain on which an iterate of f is conjuated to an irrationnal rotation on a disk. Developing an idea of Ghys, Herman proved in [Her85] a general extension theorem for holomorphic maps having an invariant annulus with rotation number in $\mathcal{H}$. Among corollaries he obtained:

Theorem (Herman)**.**

(1) *For all rational map f with degree ≥ 2 and having a Siegel disk Δ of period one and with rotation number in $\mathcal{H}$, f can not be injective in a neighborhood of $\partial\Delta$.*

(2) *For all unicritical polynomial $P(z) = z^d + c$ having a Siegel disk Δ of period one and with rotation number in $\mathcal{H}$, f has a critical point on $\partial\Delta$.*

Remark. (1) means that either f has a critical point on $\partial\Delta$ or the restriction of f to $\partial\Delta$ is non-injective (or both). [1]

The boundary $\partial\Delta$ of a Siegel disk of a rational map f of degree ≥ 2 is locally connected if and only if it is a Jordan curve (see [Mil06], Lemma 18.7). In this case, the restriction of f to $\partial\Delta$ is always injective. Thus if moreover the rotation number is in $\mathcal{H}$, then part (1) of Herman's theorem implies that there is a critical point on $\partial\Delta$. It is not known if there exist Siegel disks with non locally connected boundaries for polynomials or rational maps.

Working towards a generalization of (2) to higher degree polynomials, Rogers proved the following in [Rog98].

Theorem (Rogers)**.** *If f is a polynomial with a Siegel disk Δ of period one and rotation number in $\mathcal{H}$ then there is a critical point in the filled Siegel disk.*

---

[1] The following equivalence holds and is elementary: let K be a compact subset of the domain of definition of a holomorphic map f. Then there is no neighborhood of K on which f is injective $\iff$ either f has a critical point on K or the restriction of f to K is non-injective (or both).

The *filled Siegel disk* is defined as the union of $\partial\Delta$ and all bounded connected components of $\mathbb{C}\setminus\overline{\Delta}$. The critical point in the theorem above cannot be in the Siegel disk, thus it must be either in $\partial\Delta$ or in another bounded component of $\mathbb{C}\setminus\partial\Delta$. It is not known whether there can be such components for polynomials or rational maps.[2]

With a different approach than Herman (attacking the boundary form the inside instead of the outside, and using the Schwarzian derivative), Graczyk and Świątek proved in [GŚ03]:

Theorem (Graczyk and Świątek)**.** *If a Siegel disk has a bounded type rotation number and is compactly contained in the domain of definition of the map, then its boundary contains a critical point.*

In particular for rational maps (including polynomials) of degree ≥ 2, there is always a critical point on the boundary of bounded type Siegel disks. For polynomials, it also follows from quasiconformal surgery techniques [Zha08], which prove more:

Theorem (Zhang)**.** *The boundary of a Siegel disk of a rational map of degree* ≥ 2 *with bounded type rotation number is a Jordan curve. Thus it contains at least one critical point by Herman's theorem and the remarks that follow.*

One could conjecture that boundaries of Siegel disks of rational maps contain a critical point as soon as the rotation number is in $\mathcal{H}$. We make a small step towards this conjecture.

Theorem 1 (Main theorem)**.** *For all polynomials with two critical points, and a Siegel disk* Δ *of arbitrary period, and rotation number satisfying Herman's condition then there is an element in the cycle of* Δ *whose boundary contains a critical point.*

Our proof is a refinement on Herman's proof and uses two theorems of Mañé and the separation theorem of Goldberg Milnor Poirier Kiwi.

2. Unlikely components?

Given a bounded subset X of $\mathbb{C}$, let the filling $\operatorname{fill}(X)$ be the complement of the unbounded connected component of the complement of X: this is also the union of X and all bounded components of $\mathbb{C}\setminus X$. A bounded subset of $\mathbb{C}$ is called full if it is equal to its filling. Let

$$\widehat{X} = \operatorname{fill}(\overline{X})$$

denote the filling of the closure of X. It is a compact subset of $\mathbb{C}$. The bounded components of $\mathbb{C}\setminus\overline{X}$ will be called components *shaded* by X. For instance, we will consider $\widehat{\Delta}$ where Δ is the Siegel disk of a polynomial of degree ≥ 2. The components shaded by Δ we will be called *unlikely components.* So $\widehat{\Delta}$ is the disjoint union of Δ, $\partial\Delta$ and the unlikely components.

Remark. If Δ is a Jordan domain then there is no unlikely component. There are many polynomials and rational maps where it is known that the Siegel disk boundary is a Jordan curve. There are no polynomial or rational maps where it is known not to be. It may very well be the case that all Siegel disks of all polynomials (or rational maps) are always Jordan domains: we may be fighting against chimaeras. However, this is still unproven and seems out of reach today.

---

[2] If there is such a component, then $\partial\Delta$ cannot be locally connected (it is even worse, at least for polynomials, see Roger's theorem mentioned in section 3).

3. Remarks

Let us recall a few facts that we will not use in the present article, but which are related to the discussion.



Figure 1: Example of a simply connected open set in gray, shading a component.

On Figure 1 is an example of a simply connected open set whose closure shades a component. It cannot be a polynomial Siegel disk because the shaded component has a boundary that is not contained in the boundary of the unbounded component, contrarily to a polynomial Siegel disk. In fact it cannot be a rational Siegel disk because it has a boundary that is locally connected but not a Jordan curve.



Figure 2: Other examples.

Other examples are given on Figure 2. They cannot be polynomial Siegel disks either, yet they will serve as good illustrations for our explanations.

Rogers proved the following theorem in [Rog98]: If the polynomial f has a Siegel disk G of period one and with rotation number in $\mathcal{H}$, then either the boundary B of G contains a critical point or B is an indecomposable continuum with three properties: (1) B has at least three complementary domains, and B is the boundary of each of them, (2) each bounded complementary domain of B is a component of the grand orbit of G and so a bounded component of the Fatou set, and (3) one of the bounded complementary domains of B contains a critical point.

A *continuum* is a non-empty compact connected metric space. A continuum is *indecomposable* if it cannot be written as the union of two closed connected proper subsets. An indecomposable continuum is never locally connected. An indecomposable continuum in the plane which is the common boundary of at least three complementary domains is called a *Lakes of Wada continuum.*

There are examples of holomorphic maps defined on a simply connected open subset U of $\mathbb{C}$ with a Siegel disk compactly contained in U and whose boundary is not locally connected (for instance a pseudocircle, see [Ché11]), but no known examples with an unlikely component.

Some of the proofs in the present article are close relatives of Rogers', some will introduce new tools.

4. PROOF

Recall that a Siegel disk of a rational map is a component of the Fatou set. Recall that for a rational map, the image of a Fatou component is a Fatou component.

For $k \geq 0$, let

$$\Delta^k = P^k(\Delta)$$

so that the orbit of Δ is $\Delta = \Delta^0 \mapsto \Delta^1 \mapsto \cdots \mapsto \Delta^{p-1} \mapsto \Delta^p = \Delta$.

Let J denote the Julia set of P. Let A be the basin of attraction of infinity for P. Recall that $\partial A = J$ and that the boundary of a Siegel disk is contained in J. Thus

$$\partial\Delta^k \subset \overline{A}.$$

Recall that A is connected and unbounded. From this fact and from $\partial A = J$ it follows that any non-empty bounded and connected open subset of $\mathbb{C}$ whose boundary is contained in J cannot contain points in J and thus is a Fatou component.

Lemma 2. *Any bounded component of* $\mathbb{C} \setminus \partial\Delta^k$ *is a Fatou component.*

Proof. It follows from the above discussion, since the component has its boundary contained in $\partial\Delta^k \subset J$. □

Since Δ^k and Δ^{k+1} are Fatou components,

$$P(\partial\Delta^k) = \partial\Delta^{k+1}.$$

Recall that for a polynomial, A is completely invariant so the image of a bounded Fatou components is a bounded Fatou component.

Lemma 3. *The image of a bounded component of* $\mathbb{C} \setminus \partial\Delta^k$ *is a bounded component of* $\mathbb{C} \setminus \partial\Delta^{k+1}$*. In particular:*

$$P(\widehat{\Delta^k}) \subset \widehat{\Delta^{k+1}}.$$

Proof. Consider a bounded component U of $\mathbb{C} \setminus \partial\Delta^k$. Since $\partial U \subset \partial\Delta^k$, we have $P(\partial U) \subset \partial\Delta^{k+1}$. Now $\partial P(U) \subset P(\partial U)$ because P is open. Also, $P(U) \cap \partial\Delta^{k+1} = \varnothing$ because $P(U)$ is a Fatou component thus disjoint from $J \supset \partial\Delta^{k+1}$. Thus $P(U)$ is a connected non-empty subset of $\mathbb{C} \setminus \partial\Delta^{k+1}$ whose boundary is contained in $\partial\Delta^{k+1}$. It is therefore a component of $\mathbb{C} \setminus \partial\Delta^{k+1}$. □

Lemma 4. *Let X be a bounded subset of $\mathbb{C}$. If X is connected, then* $\operatorname{fill}(X)$ *and* $\widehat{X}$ *are connected.*

Proof. Assume by contradiction that $\operatorname{fill}(X)$ is not connected. There exists then a partition of X by two subset relatively open A and B. In particular every point in one of these sets is at positive distance from the other. Recall that $\operatorname{fill}(X)$ is the union of X and the bounded components of $\mathbb{C} \setminus X$. Each of these sets is connected thus is contained in exactly one of A and B. We may relabel them so that $X \subset A$. As $B \neq \varnothing$, there exists a bounded component M of $\mathbb{C} \setminus X$ such that $M \subset B$. As $\mathbb{C}$ is connected, there exists z in the boundary of M in $\mathbb{C}$. Either

(1) $z \in X$: then $z \in A$ thus is at positive distance from B thus from M, so it cannot belong to the boundary of M in $\mathbb{C}$. This is a contradiction.
(2) $z \notin X$: Then $M \cup \{z\}$ is a connected subset of $\mathbb{C} \setminus X$ and M a connected component thereof, so $z \in M$. Thus $z \in B$. Thus z is at positive distance from X. Thus there is some $\varepsilon > 0$ such that $B(z, \varepsilon) \cap X = \varnothing$. Thus $B(z, \varepsilon) \cup M$ is a connected subset of $\mathbb{C} \setminus X$ thus $B(z, \varepsilon) \subset M$ thus z cannot be in the boundary of M in $\mathbb{C}$. Contradiction.

Thus $\operatorname{fill}(X)$ is connected. Now if X is bounded and connected then $\overline{X}$ is bounded and connected and thus $\operatorname{fill}(\overline{X}) = \widehat{X}$ is connected. □

Thus $\widehat{\Delta^k}$ is connected. It is also contained in $P^{-1}(\widehat{\Delta^{k+1}})$ by Lemma 3.

Definition 5. *Let* $\widetilde{\Delta^k}$ *be the connected component of* $P^{-1}(\widehat{\Delta^{k+1}})$ *that contains* $\widehat{\Delta^k}$.

Let n_c the number of critical points of P and θ the rotation number of Δ. Let n_0 be the number of critical points of P belonging to $\bigcup_{k \geq 0} \widetilde{\Delta^k}$. Then $0 \leq n_0 \leq n_c$.

Theorem 2. *Let* $B = \bigcup_{k \geq 0} \partial \Delta^k$.

(1) (Herman) If $n_0 = 0$ then $\theta \notin \mathcal{H}$.
(2) If $n_0 = n_c$ then there is a critical point on B.
(3) If $n_c = 2$ and $n_0 = 1$ then the latter critical point must belong to B.

The really new part is case (3). Theorem 2 implies our main theorem, since it covers all cases when $n_c = 2$.

Complement. *If $n_0 = 0$ then there is no unlikely component.*

The proof of (1) uses classical tools of one variable complex analysis: Riemann's uniformization theorem, Schwarz's reflection principle, Caratheodory's kernel theorem. A key step is to prove that conjugating P^p by the uniformization of the complement of $\widehat{\Delta}$ to $\mathbb{C} \setminus \overline{\mathbb{D}}$ gives a circle diffeomorphism with rotation number θ.

The proof of (2) uses Sullivan's non-wandering theorem, the classification of periodic components of the Fatou set, and Mañé's theorem stating that $\partial\Delta$ is contained in the ω-limit set of a *recurrent* critical point.

The proof of (3) and the complement to (1) use the fact, already remarked by Rogers in [Rog98], that the shaded components of $\partial\Delta$ all eventually map to Δ under iteration of P. This fact is proved using the Goldberg-Milnor-Poirier-Kiwi separation theorem and Sullivan's non-wandering theorem (and indirectly the classification of periodic Fatou components via the separation theorem).

The proof of (3) will use again the classification periodic of Fatou components, and again a conjugacy of P^p by the conformal map from $\mathbb{C} \setminus \widehat{\Delta}$ to $\mathbb{C} \setminus \overline{\mathbb{D}}$, but this time giving a degree $d > 1$ cover of the circle instead of a bijection, and another theorem of Mañé, about hyperbolicity in one dimensional real dynamics, general facts about connected sets and connected components, and a little bit about plane topology and ramified covers.

4.1. Backward image of continua.

Proposition 6.

a) If $K \subset \widehat{\mathbb{C}}$ is a continuum and F a non-constant rational map then $F^{-1}(K)$ consists in finitely many connected components, all of which are mapped surjectively to K by F.

b) There exists a neighborhood V of K such that every connected component of $F^{-1}(V)$ contains at most one connected component of $F^{-1}(K)$.

Proof.

a) Let V_ε be the set of points at spherical distance at most ε from K. Then V_ε is connected and $\bigcap_{\varepsilon>0} \overline{V}_\varepsilon = K$. The map F is a finite degree ramified cover from $\mathbb{S}$ to $\mathbb{S}$. Thus F is a finite degree ramified cover from each connected component of $F^{-1}(V_\varepsilon)$ to V_ε. In particular each component of $F^{-1}(V_\varepsilon)$ is mapped surjectively to V_ε. Let C be a connected component of $F^{-1}(K)$. Let U_ε be the connected component of $F^{-1}(V_\varepsilon)$ that contains C. The set

$$L = \bigcap_{\varepsilon>0} \overline{U}_\varepsilon$$

is a decreasing intersection of compact connected sets thus is connected. It contains C since U_ε does. Since F is continuous, $F(\overline{U}_\varepsilon) \subset \overline{V}_\varepsilon$. Thus $F(L) \subset K$. Thus L is a connected subset of $F^{-1}(K)$ so $L \subset C$. So

$$L = C.$$

Now $F(U_\varepsilon) = V_\varepsilon \supset K$. For any $z \in K$ choose a point $z_\varepsilon \in U_\varepsilon \cap F^{-1}(\{z\})$. Choose any limit $x = \lim z_{\varepsilon_n}$. Then x must belong to all $\overline{U_\varepsilon}$ thus to L. Also $F(x) = z$. We have thus proved that $K \subset F(L)$. Thus

$$K \subset F(C).$$

So $K = F(C)$ i.e. C is mapped surjectively to K. Now take any point in K: it has finitely many preimage and each connected component of $F^{-1}(V_\varepsilon)$ must contain one thus there is finitely many such components.

b) The number of components of $F^{-1}(V_\varepsilon)$ is bounded from above by the degree of F and never increases as ε decreases. This number reaches thus a mamixum for ε small enough: it is equal to some $m \geq 1$ for all $\varepsilon < \varepsilon_0$. Now let $V = V_\varepsilon$ for any $\varepsilon < \varepsilon_0$. If a component of $F^{-1}(V)$ did contain more than one connected component of $F^{-1}(K)$, they would both be contained in the same intersection L, leading to a contradiction. [3] □

In the theorem above, since there are finitely many connected components of $F^{-1}(K)$ and since they are compact, each component is at positive distance from the rest. This is also implied by (b). For instance there is a neighborhood of $\widetilde{\Delta^k}$ in which the latter is the only preimage of $\widehat{\Delta^{k+1}}$, and $\widetilde{\Delta^k}$ is mapped surjectively to $\widehat{\Delta^{k+1}}$.

Lemma 7. *If $\forall k$, $\widetilde{\Delta^k} = \widehat{\Delta^{k+1}}$ then $\widehat{\Delta}$ is* locally backward invariant *by P^p, i.e. there is a neighborhood of $\widehat{\Delta}$ in which the preimage of $\widehat{\Delta}$ by P^p is $\widehat{\Delta}$.*

---

[3] This is related to the notion of *confluent* maps that we found in [Rog98]: A continuous map $f : X \to Y$ between metric spaces is confluent if each component of the inverse image of each continuum is mapped onto that continuum. Open maps between compact sets are confluent. Thus rational maps are confluent.

Proof. Without assumptions $P^p(\widehat{\Delta}) \subset \widehat{\Delta}$, i.e. $\widehat{\Delta} \subset P^{-p}(\widehat{\Delta})$. Any connected subset C of $P^{-p}(\widehat{\Delta})$ containing a point in $\widehat{\Delta}$ has connected images $P^k(C)$ that contain a point $\widehat{\Delta^k}$. By induction descending from $k = p$ to $k = 0$, it follows from the assumptions that $P^k(C) \subset \widehat{\Delta^k}$. Thus $C \subset \widehat{\Delta}$. Thus $\widehat{\Delta}$ is a connected component of $P^{-p}(\widehat{\Delta})$. By the discussion above, the union of the other components is at positive distance from it. □

The converse holds (but will not be used in this article). Note also that under the assumptions of the lemma, $P^p(\widehat{\Delta}) = \widehat{\Delta}$ so that $\widehat{\Delta}$ is in fact locally totally invariant.

Consider a non-empty connected compact subset K of $\mathbb{C}$ that is also full. Then K has a basis of connected simply connected open neighborhoods in $\mathbb{C}$; this can be proven for instance by using the conformal map ϕ from $\mathbb{C} \setminus K$ to $\mathbb{C} \setminus \overline{\mathbb{D}}$ when K is not a single point, and taking the neighborhoods $\mathbb{C} \setminus \phi^{-1}(\mathbb{C} \setminus B(0, 1+\varepsilon))$.

Lemma 8. *If $\widetilde{\Delta^k}$ does not contain a critical point then $\widetilde{\Delta^k} = \widehat{\Delta^k}$ i.e. $\widehat{\Delta^k}$ is a connected component of $P^{-1}(\widehat{\Delta^{k+1}})$. Moreover, P is a bijection from $\widehat{\Delta^k}$ to $\widehat{\Delta^{k+1}}$.*

Proof. Let us consider a connected simply connected open set V containing $\widehat{\Delta^{k+1}}$ and contained in the neighborhood of Proposition 6.b) and containing no critical point. By the discussion above, this exists. Let U be the connected component of $P^{-1}(V)$ containing $\widehat{\Delta^k}$. Then $P : U \to V$ is a cover (it has no ramification) over a simply connected set, thus is an isomorphism. In particular, P is *injective* on $\widetilde{\Delta^k}$. Since $P(\partial\Delta^k) = \partial\Delta^{k+1}$, this injectivity implies that $U \cap P^{-1}(\partial\Delta^{k+1}) = \partial\Delta^k$. Let W be a bounded component of $\mathbb{C} \setminus \partial\Delta^{k+1}$ and W' be a connected component of $P^{-1}(W) \cap U$. Then $\partial W \subset \partial\Delta^{k+1}$ thus $\partial W' \subset \partial\Delta^k$. Hence W' is contained in $\widehat{\Delta^k}$. □

4.2. **Proof of** (1)**.** It is the same as Herman's proof (that generalizes a proof of Ghys).

The composition P^p maps $\widehat{\Delta}$ in itself. By Lemma 8, and the assumption $n_0 = 0$, $\widetilde{\Delta^k} = \widehat{\Delta^k}$ for all k. Thus by Lemma 7, $\widehat{\Delta}$ is a connected component of $P^{-p}(\widehat{\Delta})$ and the rest of $P^{-p}(\widehat{\Delta})$ is at positive distance from it. Consider the unique conformal map $\phi : \mathbb{C} \setminus \widehat{\Delta} \to \mathbb{C} \setminus \overline{\mathbb{D}}$ such that $\phi(z)/z$ has a positive limit at ∞. Then a point z varying in $\mathbb{C} \setminus \widehat{\Delta}$ tends to $\widehat{\Delta}$ (for the euclidean distance on $\mathbb{C}$) if and only if $\phi(z)$ tends to $\partial\mathbb{D}$. The composition $f(z) = \phi \circ P^p \circ \phi^{-1}$ is defined on $\phi^{-1}(\mathbb{C} \setminus P^{-p}(\widehat{\Delta}))$. This set contains no point in $\overline{\mathbb{D}}$ but contains an annulus $1 < |z| < 1 + \varepsilon$ and the distance of $f(z)$ to $\partial\mathbb{D}$ tends to 0 as $z \longrightarrow \partial\mathbb{D}$. By Schwarz's reflection principle, f extends to a holomorphic map $\widetilde{f}$ defined in a neighborhood of $\partial\mathbb{D}$ and mapping $\partial\mathbb{D}$ to itself.

Lemma 9. *The restriction of $\widetilde{f}$ to $\partial\mathbb{D}$ is an analytic sense preserving diffeomorphism with rotation number θ.*

Proof. We saw that $\widetilde{f}$ is analytic. Since $\widetilde{f} = f$ for $|z| > 1$, the annulus of equation $1 < |z| < 1 + \varepsilon$ is mapped in $\mathbb{C} \setminus \overline{\mathbb{D}}$ by $\widetilde{f}$. So first there can be no critical point of $\widetilde{f}$ on $\partial\mathbb{D}$, whence f is a local diffeomorphism. Second $\widetilde{f}$ preserves the orientation on $\partial\mathbb{D}$. To prove that $\widetilde{f}$ has degree one[4] on $\partial\mathbb{D}$ and has the same rotation number as the Siegel disk, Herman's trick is to do the same construction as above but replacing $\widehat{\Delta}$ by the sub-disk $\Delta_r = \psi(\overline{B}(0, r))$ where $r < 1$ and $\psi : \mathbb{D} \to \Delta$ is a conformal map sending 0 to the center of the Siegel disk, i.e. to the periodic point.

---

[4] Being orientation preserving and without critical point is not enough to ensure bijectivity, think of the angle doubling map.

Recall that the map ψ is also a conjugacy from the rotation of angle $2\pi\theta$ to P^p. Let ϕ_r be the unique conformal map from $\mathbb{C} \setminus \Delta_r \to \mathbb{C} \setminus \overline{\mathbb{D}}$ such that $\phi_r(z)/z$ has a positive limit at ∞ and let $f_r = \phi_r \circ P^p \circ \phi_r^{-1}$. Since the boundary of Δ_r is a Jordan curve (better: it is an analytic Jordan curve), the map ϕ_r extends to a homeomorphism from $\partial\Delta_r$ to $\partial\mathbb{D}$. The map $\phi_r \circ \psi$ conjugates the rotation of angle $2\pi\theta$ on the circle of radius r to f_r on $\partial\mathbb{D}$. Thus *the rotation number of* f_r *is equal to* θ. The domain $\mathbb{C} \setminus \Delta_r$ tends to $\mathbb{C} \setminus \widehat{\Delta}$ in the sense of Caratheodory. Thus as $r \longrightarrow 1$, $\phi_r \longrightarrow \phi$ uniformly on compact subsets of $\mathbb{C} \setminus \widehat{\Delta}$ and $\phi_r^{-1} \longrightarrow \phi^{-1}$ uniformly on compact subsets of $\mathbb{C} \setminus \overline{\mathbb{D}}$. Let us prove the following claim: apart from Δ itself, the components of $P^{-p}(\Delta)$ are at positive distance from $\widehat{\Delta}$. First recall that $P^{-p}(\widehat{\Delta}) \setminus \widehat{\Delta}$ is at positive distance from $\widehat{\Delta}$. So we must prove that $P^{-p}(\Delta) \cap \widehat{\Delta} = \Delta$. This follows from the injectivity of P on $\widehat{\Delta^k}$ given by Lemma 8. We proved the claim. Now, since $\phi_r^{-1} \longrightarrow \phi^{-1}$ uniformly on compact sets, the image by ϕ_r^{-1} of the annulus of equation $1 < |z| < 1 + \varepsilon/2$ is disjoint from the components of $P^{-p}(\Delta)$ that are different from Δ. Thus the domains of definition of f_r contain a common annulus $1 < |z| < 1 + \varepsilon/2$ for r close enough to 1. Thus the domains of definition of $\widetilde{f}_r$ contain a common annulus containing $\partial\mathbb{D}$ for r close enough to 1. On the boundary of this annulus, $\widetilde{f}_r$ converges uniformly to $\widetilde{f}$. This implies by the maximum principle that $\widetilde{f}_r$ converges uniformly to $\widetilde{f}$ on the annulus, so in particular on $\partial\mathbb{D}$. As a uniform limit of sense preserving homeomorphisms of $\widetilde{f}_r : \partial\mathbb{D} \to \partial\mathbb{D}$ with rotation number θ, $\widetilde{f}$ has degree one and rotation number θ. □

Now if θ did belong to $\mathcal{H}$ then $\widetilde{f}$ would be analytically linearizable on $\partial\mathbb{D}$. The linearizing map would have an injective holomorphic extension to a neighborhood of $\partial\mathbb{D}$, and by analytic continuation, this extension would still conjugate $\widetilde{f}$ to a rotation in a neighborhood of $\partial\mathbb{D}$. Recall that $\widetilde{f}(z) = f(z)$ for $|z| > 1$ thus f would have a stable domain containing an annulus of the form $1 < |z| < 1 + \varepsilon'$. Thus P^p would have a stable domain containing a neighborhood of $\widehat{\Delta}$, thus a neighborhood of $\partial\Delta$. This contradict the fact that $\partial\Delta$ is contained in the Julia set.

4.3. **Proof of** (2).

Denote $\omega(z)$ the ω-limit set of z, i.e. the set of points of accumulation of the sequence $P^n(z)$.

Consider a critical point c that belongs to the Fatou set of P. By Sullivan's non-wandering theorem and Fatou's classification of periodic components, $\omega(c)$ is either equal to an attracting cycle, a parabolic cycle, or a cycle of invariant curves in Siegel disks. The intersection $\omega(c) \cap J$ is therefore finite or empty.

Another theorem of Fatou says that $\partial\Delta \subset \bigcup_c \omega(c)$, the union being over all critical points. The set $\partial\Delta$ is contained in J and contains infinitely many points.

Recall that $\widehat{\Delta^k}$ is the disjoint union of $\partial\Delta^k$ and of the bounded components of $\mathbb{C}\setminus\partial\Delta^k$, and that the latter are Fatou components. Since we are in the case $n_0 = n_c$, all critical values of P belong to some $\widehat{\Delta^k}$. If no critical value were on B, they would all belong to the Fatou set, thus $J \cap \left(\bigcup_c \omega(c)\right)$ would be finite contradicting the fact that it contains the infinite set $\partial\Delta$.

To prove that there is not only a critical value, but also a critical point on B, we must use the following theorem of Mañé [Mañ93], much harder than Fatou's: there exists a *recurrent* critical point c such that $\partial\Delta \subset \omega(c)$. As above, $P(c)$ must belong to B. Since $P(B) = B$, we have $\omega(c) = \omega(P(c)) \subset B$. Since $c \in \omega(c)$, we have $c \in B$.

Remark. We did not need the following, but in fact according to Roger's lemma below, every unlikely component is eventually mapped to a Δ^k. Thus if $P(c)$ belongs to an unlikely component, then $\omega(c)$ is a cycle of invariant curves in $\bigcup_k \Delta^k$, and thus does not intersect J at all.

4.4. **Unlikely components eventually map to the Siegel disk.** To prove (3) and the complement to (1), we need the following lemma. Recall that a component shaded by $\partial\Delta^k$ is a Fatou component and is mapped to a component shaded by $\partial\Delta^{k+1}$.

Lemma 10 (Rogers, Theorem 3.3 in [Rog98])**.** *All components shaded by* $\partial\Delta^k$ *eventually map under iteration of* P *to some* Δ^l *(thus eventually maps to any* Δ^l*).*

Proof. This is almost the same proof as in [Rog98]. The Goldberg-Milnor-Poirier-Kiwi separation theorem [Kiw00] states that there exists $m > 0$ such that the union L of the set of external rays fixed by P^m (there are finitely many of them) union their endpoints (they land) cut the plane into regions, each of them containing at most one periodic Fatou component or cremer point (and never both of them).

Let U be a components shaded by $\partial\Delta^k$. By Sullivan's non-wandering theorem, it is eventually mapped to a periodic component $V = P^n(U)$. Such a component is by Fatou's classification either an attracting component, a parabolic component, or a Siegel component. If V were not one of the Δ^l, then V would be separated by L from all the Δ^l. In particular, the intersection of ∂V with $\partial\Delta^l$ would be at most one point. However, $\partial V \subset B$ and ∂V is infinite. Contradiction. □

4.5. **Proof of the complement to** (1)**.** If $n_0 = 0$ then we saw that the map P is a bijection from $\widehat{\Delta^k}$ to $\widehat{\Delta^{k+1}}$. Thus Δ^{k+1} has only one preimage by P in $\widehat{\Delta^k}$, namely Δ^k. Any component V shaded by Δ^m has its orbit contained in $\bigcup_k \widehat{\Delta^k}$, and eventually maps to some Δ^l by Rogers' lemma. Thus $V = \Delta^m$. There is thus only one component shaded by $\partial\Delta^m$: there is no unlikely component.

4.6. **More tools.** Recall that $\widetilde{\Delta^k}$ is the connected component of $P^{-1}(\widehat{\Delta^{k+1}})$ containing $\widehat{\Delta^k}$. We can choose a small enough connected and simply connected open set V_k containing $\widehat{\Delta^k}$ so that there is no critical value of P in $V_k \setminus \widehat{\Delta^k}$. By Proposition 6.b), we can choose V_k so that moreover, letting U_k be the connected component of $P^{-1}(V_{k+1})$ that contains $\widehat{\Delta^k}$, we have $U_k \cap P^{-1}(\widehat{\Delta^{k+1}}) = \widetilde{\Delta^k}$ i.e. U_k contains $\widetilde{\Delta^k}$ but does not contain the other preimages of points of $\widehat{\Delta^{k+1}}$ if there are some. Let us also set $U_{k+p} = U_k$ and $V_{k+p} = V_k$ for all $k \geq 0$ and $n \geq 0$. The map P_k is a finite degree ramified cover from U_k to V_{k+1}. The set U_k is simply connected.

Note that there is no a priori inclusion between U_k and V_k.

By the separation theorem, there is a system of external rays and their endpoints that separate the different Δ^k. The external rays are contained in the basin of infinity thus they do not intersect $\widehat{\Delta^k}$. Thus for $i \neq j \bmod p$, $\widehat{\Delta^i} \cap \widehat{\Delta^j}$ is either empty or consists of a single landing point of a periodic external ray.

The following lemma is elementary:

Lemma 11. *If* A *and* B *are bounded subsets of* $\mathbb{C}$ *then*

$$A \subset \operatorname{fill}(B) \iff \operatorname{fill}(A) \subset \operatorname{fill}(B).$$

Proof.
$\Longleftarrow$: immediate since $A \subset \operatorname{fill}(A)$.
$\Longrightarrow$: when $A \subset \operatorname{fill}(B)$, the unbounded component of $\mathbb{C} \setminus B$ is disjoint from A thus it is an unbounded connected subset of $\mathbb{C} \setminus A$ thus it is contained in the unbounded component of $\mathbb{C} \setminus A$, i.e. $\operatorname{fill}(A) \subset \operatorname{fill}(B)$. □

Corollary 12. *If A and B are bounded subsets of $\mathbb{C}$ then*

$$A \subset \widehat{B} \iff \overline{A} \subset \widehat{B} \iff \widehat{A} \subset \widehat{B}.$$

The first equivalence holds because $\widehat{B}$ is closed, the second by applying Lemma 11 to $\overline{A}$ and $\overline{B}$.

Lemma 13. *For all* compact *subset K of $\mathbb{C}$, $P^{-1}(\mathrm{fill}(K)) = \mathrm{fill}(P^{-1}(K))$.*

Proof. Let U be the unbounded component of $\mathbb{C} \setminus K$, so that $\mathrm{fill}(K) = \mathbb{C} \setminus U$. Let V be the unbounded component of $\mathbb{C} \setminus P^{-1}(K)$. The lemma is equivalent to $V = P^{-1}(U)$. As V is connected, $P(V)$ is connected and disjoint from K and undbounded thus $P(V) \subset U$, i.e. $V \subset P^{-1}(U)$. Since P is a finite degree ramified cover and U is open, its restriction to $P^{-1}(U)$ is a finite degree ramified cover over U. Each component of $P^{-1}(U)$ will map surjectively to U, and thus be unbounded, so there is exactly one component of $P^{-1}(U)$. Thus $P^{-1}(U)$ is unbounded and connected and it does not intersect $P^{-1}(K)$ (as U does not intersect K). Thus $P^{-1}(U) \subset V$. □

We do not know if Lemma 13 still holds if we replace the compact K by a bounded subset X. However:

Corollary 14. *For all bounded subset X of $\mathbb{C}$, $P^{-1}(\widehat{X}) = \widehat{P^{-1}(X)}$.*

Proof. By definition, $\widehat{P^{-1}(X)} = \mathrm{fill}(\overline{P^{-1}(X)})$. Next, $\overline{P^{-1}(X)} = P^{-1}(\overline{X})$ ($\subset$ because P is continuous, $\supset$ because P is open). Now apply Lemma 13 with $K = \overline{X}$. □

The following lemmas aim at proving Proposition 17 below.

Lemma 15. *If X and Y are bounded subsets of $\mathbb{C}$ and $\partial Y \subset X$ then $\overline{Y} \subset \mathrm{fill}(X)$.*

Proof. Let A be the unbounded component of $\mathbb{C} \setminus X$: if $Y \not\subset \mathrm{fill}(X)$ then $Y \cap A \neq \varnothing$. As Y is bounded, $A \not\subset Y$ and as A is connected there exists $z \in A$ in the boundary of $Y \cap A$ relative to A. This point is also in the boundary of Y relative to $\mathbb{C}$, thus in X by asumption. Thus $z \in X \cap A = \varnothing$, which is a contradiction. We have thus proved that $Y \subset \mathrm{fill}(X)$. Since $\overline{Y} = Y \cup \partial Y$, and $\partial Y \subset X \subset \mathrm{fill}(X)$, we get $\overline{Y} \subset \mathrm{fill}(X)$. □

Lemma 16. *The connected components of a full bounded set are full (and bounded).*

Proof. Let X be a full bounded set, so $\mathrm{fill}(X) = X$. Let C be a component of X. By Lemma 4, $\mathrm{fill}(C)$ is connected. Since $C \subset X$, Lemma 11 gives $\mathrm{fill}(C) \subset \mathrm{fill}(X)$, i.e. $\mathrm{fill}(C) \subset X$. Thus $\mathrm{fill}(C)$ is a connected subset of X containing the connected component C of X thus $\mathrm{fill}(C) = C$, i.e. C is full. □

Proposition 17. *Let X be a bounded subset of $\mathbb{C}$.*

1. *For all connected component C of $\mathrm{fill}(X)$,*

$$C = \mathrm{fill}(X \cap C).$$

2. *For all connected component C of $\widehat{X}$,*

$$C = \widehat{\overline{X} \cap C}.$$

Proof. (1) Let us prove the two inclusions $C \supset \mathrm{fill}(X \cap C)$ and $C \subset \mathrm{fill}(X \cap C)$.
$\supset$: $X \cap C \subset C \subset \mathrm{fill}(C)$ thus by Lemma 11, $\mathrm{fill}(X \cap C) \subset \mathrm{fill}(C)$. The set $\mathrm{fill}(X)$ is full so by Lemma 16, $\mathrm{fill}(C) = C$.
$\subset$: Let $z \in C$. Either

- $z \in X$: Then $z \in X \cap C \subset \mathrm{fill}(X \cap C)$.

- $z \notin X$: Let A the bounded component of $\mathbb{C} \setminus X$ to which z belongs. Then $A \subset C$. Let B the unbounded component of $\mathbb{C} \setminus X$. Since A and B are distinct connected components of $\mathbb{C} \setminus X$, there exists an open subset U of $\mathbb{C}$ such that $A \subset U$, $B \cap U = \varnothing$, and $\partial U \subset X$. Let V be the connected component of U that contains A. Then $\partial V \subset \partial U$ thus $\partial V \subset X$. By Lemma 15, $\overline{V} \subset \operatorname{fill}(X)$. Since V is connected, $\overline{V}$ is connected. Since C is a connected component of $\operatorname{fill}(X)$, $\overline{V} \subset C$. Therefore $\partial V \subset X \cap C$. Moreover $V \cap B \subset U \cap B = \varnothing$ thus $V \cap B = \varnothing$. Therefore, z is contained in a bounded connected component of $\mathbb{C} \setminus (X \cap C)$: $z \in \operatorname{fill}(X \cap C)$.

(2) comes from (1) applied to $\overline{X}$, since $\widehat{X} = \operatorname{fill}(\overline{X})$, and from the fact that C is closed, as a connected component of a closed set, thus $\overline{X} \cap C$ is closed thus $\operatorname{fill}(\overline{X} \cap C) = \widehat{\overline{X} \cap C}$. □

Remark. Proposition 17, item (2) is not true if we replace $\overline{X}$ by X, as the following example shows (see figure 3): X is the union of the three (closed) segments drawing a polygonal line with vertices 0, $-i$, $1-i$, 1, and of the horizontal segments $[0,1] + i/2^n$ for $n > 0$, and C is the convex hull of the four aforementioned vertices.



Figure 3: An example of bounded set X and connected component C of $\widehat{X}$ where $C \neq \widehat{X \cap C}$.

4.7. **Proof of** (3)**.** Now let us deal with the case where $n_0 = 1$: there is only one critical point in the union of the $\widetilde{\Delta^k}$. Call it c'. Without loss of generality, we may assume that $c' \in \operatorname{fill}(\Delta)$. In section 4.6 we defined simply connected open sets V_k and U_k containing $\widehat{\Delta^k}$, and that there is no critical value of P in $V^k \setminus \widehat{\Delta^k}$ and P is a ramified cover from U_k to V_{k+1} and $U_{k+p} = U_k$ and $V_{k+p} = V_k$.

First case: $P(c') \in \partial\Delta^1$. The critical value $P(c') \in V_1$ has only one preimage in U_0. Since there must be at least one preimage on $\partial\Delta$ (recall that $P(\partial\Delta^k) = \partial\Delta^{k+1}$), this implies that $c' \in \partial\Delta$.

Second case: $P(c') \notin \partial\Delta^1$. We will get a contradiction according to the following plan: we will first prove that $\widehat{\Delta}$ is locally totally invariant by P^p (see below). We will then conjugate P^p by the conformal map from $\mathbb{C} \setminus \widehat{\Delta} \to \mathbb{C} \setminus \overline{\mathbb{D}}$, like in case (1). The local total invariance implies that we get a map defined in an annulus $1 < |z| < 1 + \varepsilon$, but this time the induced circle map has degree > 1. We will then use a theorem of Mañé ensuring hyperbolicity of this circle map, provided there are no non repelling cycles on the circle. The hypothesis $n_c = 2$ of case (3) will be used to check the absence of such cycles. Once we have this hyperbolicity, this implies that the map has a polynomial-like restriction to a neighborhood of $\widehat{\Delta}$, with only one critical point. We can then apply Herman's theorem (or case (2)), and get the contradiction.

The rest of section 4.7 is devoted to the proof of the contradiction in the second case.

4.7.1. *Towards local invariance.* Recall that we work by contradiction, so that what we prove here needs not to hold in reality. As $P^p(\widehat{\Delta}) = \widehat{\Delta}$, we have $\widehat{\Delta} \subset P^{-p}(\widehat{\Delta})$. Let us prove that those points of the p-th preimage of $\widehat{\Delta}$ that are not in $\widehat{\Delta}$ are at some distance from $\widehat{\Delta}$, i.e. that $\widehat{\Delta}$ *is locally totally invariant by* P^p. For this it is sufficient to prove that $\forall k$, $\widetilde{\Delta^k} = \widehat{\Delta^k}$. (^{5})

The assumption that the critical value $P(c')$ does not belong to $\partial\Delta^1$ implies that it belongs to a component shaded by $\partial\Delta^1$. Since on one hand $P(c')$ belongs to the Fatou set and on the other hand $\widehat{\Delta^i} \cap \widehat{\Delta^j}$ is either empty or a single point in the Julia set, it follows that there is no critical value in $\widehat{\Delta^k}$ when $k \neq 1 \bmod p$. Thus by Lemma 8:

$$\text{when } k \neq 0 \bmod p,\ \widetilde{\Delta^k} = \widehat{\Delta^k}.$$

The next few sections aim to prove that it also holds when $k = 0 \bmod p$, i.e. that $\widetilde{\Delta} = \widehat{\Delta}$. We saw that $P(c')$ belongs to a component shaded by $\partial\Delta^1$. It cannot belong to Δ^1 for otherwise it would have a simple preimage in Δ but it has only one preimage in U_0, which is multiple. Thus $P(c')$ belongs to an unlikely component. This unlikely component has its orbit contained in $\bigcup_k \widehat{\Delta^k}$ and is eventually mapped to some Δ^l.

4.7.2. *Symmetries.* The map $P : U_0 \to V_1$ is a ramified cover, ramified over only one point, and V_1 is simply connected thus it is equivalent to the map $D : \mathbb{D} \to \mathbb{D}$, $z \mapsto z^d$ where d is the local degree of P at c'. Since $P(c')$ does not belong to Δ^1, it implies that Δ^1 has exactly d preimages in $\widetilde{\Delta}$. The map $D : z \mapsto z^d$ has a transitive and cyclic group of automorphisms $\simeq (\mathbb{Z}/d\mathbb{Z}, +)$: it is the group of rotations generated by $z \mapsto e^{2\pi i/d}$. Thus the ramified cover $P : U_0 \to V_1$ has a transitive group of automorphisms G that is cyclic. Let $\rho : U_0 \to U_0$ be a generator. The d preimages of Δ^1 are the sets $\rho^n(\Delta)$, $n \in \mathbb{Z}$. Let us call them *partners* and denote $\mathcal{O}$ their union:

$$\mathcal{O} = \widetilde{\Delta} \cap P^{-1}(\Delta^1) = U_0 \cap P^{-1}(\Delta^1) = \bigcup_{n\in\mathbb{Z}} \rho^n(\Delta) = \bigcup_{g\in G} g(\Delta).$$

---

[5] Indeed, the latter means that every point close enough to (i.e. in a neighborhood of) $\widehat{\Delta^k}$ but not belonging to this set is mapped to the complement of $\widehat{\Delta^{k+1}}$ by P (see the discussion in Section 4.1). By continuity of P, points close to $\widehat{\Delta^k}$ are mapped close to $\widehat{\Delta^{k+1}}$. Thus for all $m > 0$ and all k, points close enough to $\widehat{\Delta^k}$ but not belonging to this set are mapped by P^m to the complement of $\widehat{\Delta^{k+m}}$.

By Corollary 14, $P^{-1}(\widehat{\Delta^1}) = \widehat{P^{-1}(\Delta^1)}$. The set $\widetilde{\Delta}$ is thus a connected component of $\widehat{P^{-1}(\Delta^1)}$, so by Proposition 17, $\widetilde{\Delta} = \mathrm{fill}(\overline{P^{-1}(\Delta^1)} \cap \widetilde{\Delta})$. Now $P^{-1}(\Delta^1)$, the preimage by P of Δ^1, consists in finitely many components: the ones in $\widetilde{\Delta}$, whose union compose $\mathcal{O}$ and the other ones, which are disjoint from U_0, so their closure is disjoint from $\widetilde{\Delta}$. Thus $\overline{P^{-1}(\Delta^1)} \cap \widetilde{\Delta} = \overline{P^{-1}(\Delta^1) \cap \widetilde{\Delta}}$. Thus $\mathrm{fill}(\overline{P^{-1}(\Delta^1)} \cap \widetilde{\Delta}) = \mathrm{fill}(\overline{P^{-1}(\Delta^1) \cap \widetilde{\Delta}}) = \mathrm{fill}(\overline{Z}) = \widehat{Z}$ with $Z = P^{-1}(\Delta^1) \cap \widetilde{\Delta} = \mathcal{O}$. Thus

$$\widetilde{\Delta} = \widehat{\mathcal{O}}.$$

4.7.3. *Mutual shading.* Given what was said above and since $P(c') \notin \Delta^1$, the first time c' gets in $\bigcup_k \Delta^k$ is for the form $P^{1+mp}(c') \in \Delta^1$ with $m \geq 1$. Thus $P^{mp}(c') \in u(\Delta)$ for some $u \in G$ with $u \neq \mathrm{id}$. Recall $P(c')$ is contained in a component shaded by $\partial \Delta^1$. Since $mp \geq 1$, it follows that $P^{mp}(c')$ belongs to a component shaded by $\partial \Delta$. So we proved that:

$$\exists u \in G \setminus \{\mathrm{id}\} \quad u(\Delta) \subset \widehat{\Delta}.$$

In other words, Δ *shades one of its partners.*

By Corollary 12: $\widehat{u(\Delta)} \subset \widehat{\Delta}$. Since U_0 is simply connected, its complement is connected and thus for any subset X of U_0 and any $a \in G$, $\mathrm{fill}(a(X)) = a(\mathrm{fill}(X))$, and thus $\widehat{a(X)} = a(\widehat{X})$. In particular

$$u(\widehat{\Delta}) \subset \widehat{\Delta}.$$

Now let M be any element of the orbit of $\widehat{\Delta}$ under G and assume that for some $a \in G$, $a(M) \subset M$. Then $\forall g \in G$, $ga(M) \subset g(M)$. In particular $M = a^d(M) \subset a^{d-1}(M) \subset \cdots \subset a^2(M) \subset a(M) \subset M$. Thus $a(M) = M$. In particular

$$u(\widehat{\Delta}) = \widehat{\Delta}.$$

Applying $g \in G$, we get: $ga(M) = g(M)$. Since G is a cyclic group, there exists a divisor $m > 0$ of d such that, noting $H = \langle \rho^m \rangle$ the subgroup generated by ρ^m, and $G \cdot \widehat{\Delta} = \{\widehat{\Delta}, \rho(\widehat{\Delta}), \rho^2(\widehat{\Delta}), \ldots\}$, we have:

$\forall a \in G, \quad \forall M, N \in G \cdot \widehat{\Delta}$:

- $a(M) = M \iff a \in H$,
- $M \subset N \implies M = N$.

In particular,

$$|H| > 1.$$

Note that by the second point, $M \neq N \implies M \not\subset N$ *and* $N \not\subset M$.

For all $h \in H$, $h(\partial \Delta) = h(\partial \widehat{\Delta}) = \partial h(\widehat{\Delta}) = \partial \widehat{\Delta} = \partial \Delta$:

$$\forall h \in H, \quad h(\partial \Delta) = \partial \Delta.$$

For all $g \in G$, let $M = g(\Delta)$: recall that $c' \in \widetilde{\Delta} = \widehat{\Delta}$. Thus $g(c') \in g(\widehat{\Delta}) = \widehat{g(\Delta)}$. Since c' is fixed by g: $c' \in \widehat{M} \setminus \overline{M}$, i.e. the critical point c' is shaded by $g(\Delta)$.

4.7.4. *Wada lakes.* We will prove that $H = G$. Therefore, all partners will shade each other, and they will all have the same boundaries, like Wada lakes.

Assume by contradiction that $H \neq G$.

Then $\widehat{\Delta} \neq \rho(\widehat{\Delta})$ thus by the properties above, $\widehat{\Delta} \not\subset \rho(\widehat{\Delta})$ and $\rho(\widehat{\Delta}) \not\subset \widehat{\Delta}$.

As $\widehat{\Delta} \not\subset \rho(\widehat{\Delta})$ and $\widehat{\Delta} = \widehat{\partial \Delta}$ and $\rho(\widehat{\Delta}) = \widehat{\rho(\Delta)}$ this implies (see Corollary 12) that $\partial \Delta \not\subset \rho(\widehat{\Delta})$. Thus there is a point $z_0 \in \partial \Delta \setminus \rho(\widehat{\Delta})$. Since $\rho(\widehat{\Delta})$ is closed, there is a neighborhood of z_0 that does not intersect $\rho(\widehat{\Delta})$. Let $B = B(z_0, \varepsilon)$ be included in this neighborhood and in U_0 and not containing the critical point. Since $z_0 \in \partial \Delta$, B intersects Δ. Denote $b = \rho^m$, which we recall is a generator of H. Since $b \in H$, $b(\partial \Delta) = \partial \Delta$ so $b(z_0) \in \partial \Delta$ thus $b(B)$ intersects Δ too. Now choose any curve contained in Δ and going from a point z_1 in $\Delta \cap B$ to a point z_2 in $\Delta \cap h(B)$. Complete it by any curve contained in B from z_1 to z_0 and by any curve contained in $h(B)$ from z_2 to $h(z_0)$: thus we get a curve γ_0 going from z_0 to $h(z_0)$. Extend this curve by periodicity into a curve $\gamma = \gamma_0 \cdot b(\gamma_0) \cdot b^2(\gamma_0) \cdot \ldots$, we get a closed curve[6] going through z_0, $b(z_0)$, ... This curve does not contain the critical point, it does not intersect $\rho(\widehat{\Delta})$ (so in particular $\gamma \cap \rho(\partial\Delta) = \varnothing$) and for all $h \in H$ it does not intersect $\rho h(\Delta)$.

Let us prove that γ separates c' from ∞. For this let us go into the disk model of U_0: we already remarked that there is an analytic isomorphism from U_0 to $\mathbb{D}$ that sends c' to 0 and conjugates the group G to the group of rotations generated by $z \mapsto e^{2\pi i/d} z$. The curve γ_0 goes from z_0 to some $e^{2\pi i k/d} z_0 \neq z_0$. The lifted rotation angle around 0 of γ_0 is therefore $\neq 0$. The lifted rotation angle of γ is d times the former, thus $\neq 0$. Therefore the winding number of γ around 0 is $\neq 0$. So γ separates 0 from $\partial\mathbb{D}$ in the disk model. This proves the claim.

Let us sum up: using $\widehat{\Delta} \not\subset \rho(\widehat{\Delta})$ we have constructed a closed curve γ that separates c' from ∞, contains a point in $\partial\Delta$, does not intersect $\rho(\partial\Delta)$ and for all $h \in H$ does not intersect $\rho h(\Delta)$.

Similarly using $\rho(\widehat{\Delta}) \not\subset \widehat{\Delta}$ we can construct a closed curve γ' that separates c' from ∞, contains a point in $\rho(\partial\Delta)$, does not intersect $\partial\Delta$ and for all $h \in H$ does not intersect $h(\Delta)$.

We claim that the construction of γ' can be done so that moreover:

$$\gamma \cap \gamma' = \varnothing.$$

Indeed, $\gamma \subset \Delta \cup \bigcup_{h\in H} h(B)$ and $\gamma' \subset \rho(\Delta) \cup \bigcup_{h \in H} h(B')$ and $h(B)$ doet not meet $\rho(\widehat{\Delta})$ (because $h(B) \cap \rho(\widehat{\Delta}) = h(B) \cap \rho(h(\widehat{\Delta})) = h(B) \cap h(\rho(\widehat{\Delta})) = h(B \cap \rho(\widehat{\Delta})) = h(\varnothing) = \varnothing$). In particular $h(B) \cap \rho(\Delta) = \varnothing$. Similarly $h(B') \cap \Delta = \varnothing$. Of course $\Delta \cap \rho(\Delta) = \varnothing$. So the only possible intersection is in some $h(B) \cap h'(B')$ for some $h, h' \in H$. So once the starting point z_0' has been chosen in $\rho(\partial\Delta) \setminus \widehat{\Delta}$, which is non-empty, then note that $h(z_0')$ does not belong to γ because γ is disjoint from $h\rho(\widehat{\Delta}) = \rho h(\widehat{\Delta}) = \rho(\widehat{\Delta})$ and then it is enough to choose the ball $B' = B(z_0', \varepsilon')$ small enough so that $\forall h \in H$, $h(B') \cap \gamma = \varnothing$.

Let X be a bounded subset of $\mathbb{C}$. Let $cc_\infty(X)$ denote the unbounded component of $\mathbb{C} \setminus X$. The letters cc stand for complementary component. For $z \notin X$, let $cc_z(X)$ denote the component of $\mathbb{C} \setminus X$ containing z. If Z is a non empty subset of a connected component U of $\mathbb{C} \setminus X$, let $cc_Z(X)$ denote U.

Let $X = \rho(\partial\Delta)$ and $Y = \gamma$. Then X and Y are bounded, non empty, disjoint, connected. Thus at least one of them must be contained in the unbounded component of the complement of the other: $X \subset cc_\infty(Y)$ or $Y \subset cc_\infty(X)$ or both. Similarly, neither X nor Y contains c' thus, $X \subset cc_{c'}(Y)$ or $Y \subset cc_{c'}(X)$ or both. Both X and Y separate c' from ∞: $cc_{c'}(X) \neq cc_\infty(X)$ and $cc_{c'}(Y) \neq cc_\infty(Y)$. From all this, it follows that exactly one of the following holds:

- $X \subset cc_\infty(Y)$ and $Y \subset cc_{c'}(X)$
- $Y \subset cc_\infty(X)$ and $X \subset cc_{c'}(Y)$

Now, $Y = \gamma$ cannot be contained in a bounded component of the complement of $X = \rho(\partial\Delta)$, for otherwise γ would be contained in a Fatou component but γ contains a point of $\partial\Delta$. Thus it is the second line that holds:

$$\gamma \subset cc_\infty(\rho(\partial\Delta)) \text{ and } \rho(\partial\Delta) \subset cc_{c'}(\gamma).$$

Similarly

$$\gamma' \subset cc_\infty(\partial\Delta) \text{ and } \partial\Delta \subset cc_{c'}(\gamma').$$

---

[6] It is hot hard to get a *simple* closed curve, but we will not need to.

Now γ' is disjoint from γ and contains a point in $\rho(\partial\Delta)$ thus $\gamma' \subset cc_{c'}(\gamma)$. Similarly $\gamma \subset cc_{c'}(\gamma')$. This contradicts the fact that γ and γ' separate c' from ∞.

We thus proved that $H \neq G$ leads to a contradiction.

4.7.5. *Obtaining local invariance.* Therefore $H = G$. We have thus proved that $\forall g \in G$, $g(\widehat{\Delta}) = \widehat{\Delta}$. Thus

$$\forall g \in G,\ \widehat{g(\Delta)} = \widehat{\Delta}.$$

In particular $g(\Delta) \subset \widehat{\Delta}$. Thus we proved: Δ *shades all its partners* i.e.

$$\mathcal{O} \subset \widehat{\Delta}.$$

Using Corollary 12 it follows that $\widehat{\mathcal{O}} \subset \widehat{\Delta}$. Recall that $\widetilde{\Delta} = \widehat{\mathcal{O}}$. Thus we proved that $\widetilde{\Delta} \subset \widehat{\Delta}$. Since by definition $\widehat{\Delta} \subset \widetilde{\Delta}$, we proved:

$$\widetilde{\Delta} = \widehat{\Delta}.$$

This finishes the proof that $\widehat{\Delta}$ is locally totally invariant by P^p, under the asumption that $P(c') \notin \partial\Delta^1$.

4.7.6. *Analytic cover of the circle with non vanishing derivative.* Therefore, like in case (1) of Theorem 2, conjugating P^p by a conformal isomorphism $\phi : \mathbb{C}\backslash\widehat{\Delta} \to \mathbb{C}\backslash\mathbb{D}$ yields an analytic function $f = \phi \circ P^p \circ \phi^{-1}$ whose domain of definition contains an annulus of the form $1 < |z| < 1 + \varepsilon$. Again, $|f(z)|$ tends to 1 as $|z| \longrightarrow 1$ so we can do Schwarz's reflection and get an analytic extension $\widetilde{f}$ to a neighborhood of the unit circle ∂D. The difference with case (1) is that the new map is no longer a circle diffeomorphism:

Lemma 18. *The restriction of $\widetilde{f}$ to $\partial\mathbb{D}$ is a is a degree d cover of the circle, with non-vanishing derivative.*

Proof. The derivative cannot vanish and the differential must preserve the orientation of $\partial\mathbb{D}$. Those two claims hold because $\widetilde{f}(z) = f(z)$ on $1 < |z| < 1 + \varepsilon$ and the image of this annulus by f does not intersect $\mathbb{D}$. Thus $\widetilde{f}$ must be a cover of the circle. Let m be its degree. For all neighborhood U of $\partial\mathbb{D}$ there exists a neighborhood V of $\partial\mathbb{D}$ such that every point in $V \setminus \overline{\mathbb{D}}$ has exactly d preimages by f in $U \setminus \overline{\mathbb{D}}$. The map P^p is a degree d cover from U_0 to V_1 (whose critical value is in $\widehat{\Delta}$) and an isomorphism from U_k to V_{k+1} when $k \neq 0 \bmod p$. Thus we have a similar property: for all neighborhood U of $\widehat{\Delta^k}$, there exists a neighborhood V of $\widehat{\Delta^{k+1}}$ such that every point of $V \setminus \widehat{\Delta^{k+1}}$ has exactly d or 1 preimages by P^p in $U \setminus \widehat{\Delta^k}$, according to whether $k = 0 \bmod p$ or not. From this it follows that for all neighborhood U of $\widehat{\Delta}$ there exists a neighborhood V of $\widehat{\Delta}$ such that every point in $V \setminus \widehat{\Delta}$ has exactly d preimages in $U \setminus \widehat{\Delta}$. Thus $m = d$. □

4.7.7. *Absence of non repelling cycle for the circle map.* Let us work by contradiction. In the case of existence of an attracting or parabolic cycle of $\widetilde{f}$ on $\partial\mathbb{D}$, choose a point a in the cycle, let m be its period and let $\mathcal{A}$ be its immediate basin. Let $\mathcal{B} = \phi^{-1}(\mathcal{A})$. Then $\mathcal{B}$ is bounded in $\mathbb{C}$ and stable by an iterate of P^{mp}. It is therefore contained in a periodic component of the Fatou set $\mathcal{B}'$. Let us prove $\mathcal{B}' = \mathcal{B}$, even though we do not really need that fact: $\phi(\mathcal{B}')$ is bounded, connected, invariant by $\widetilde{f}$ and contains $\mathcal{A}$. It is thus contained in the immediate basin of the same attracting periodic point/parabolic petal, a.k.a. $\mathcal{A}$. Thus $\phi(\mathcal{B}') = \mathcal{A}$ thus $\mathcal{B}' = \mathcal{B}$. Since every point of $\mathcal{A}$ tends to a under iteration of $\widetilde{f}^{mp}$, every point of $\mathcal{B}$ tends to $\partial\Delta$ under iteration of P^{mp}. Thus in particular the component $\mathcal{B}$ is not part of a cycle of Siegel disks, because every point in a Siegel disk has an orbit that stays bounded away from the Julia set. It is therefore a component of an immediate attracting or parabolic basin. In particular the cycle of components $\mathcal{B}$ belongs to must contain a critical point.

Let us now use the assumption $n_c = 2$, i.e. that P only has two critical points. One of them is c' and eventually maps to Δ so its orbit does not accumulate on $\partial\Delta$. The other is contained in the cycle of $\mathcal{B}$ so its orbit accumulates on a finite set. This contradicts Fatou's theorem saying that $\partial\Delta \subset \bigcup_c \omega(c)$, the union being over all critical points.

Therefore there is no non-repelling cycle for $\widetilde{f}$ on $\partial\mathbb{D}$.

Remark. If there is $n_c > 2$ critical points then we were not able to get a contradiction, because a third critical point could play the role in Fatou's theorem. All we can say is that if there is a non-repelling cycle for $\widetilde{f}$ on $\partial\mathbb{D}$ then there must be a parabolic cycle on $\partial\Delta$. Indeed in $\mathcal{B}$ every point tends to $\partial\Delta$ under iteration of P^{mp} but also to the attracting or parabolic cycle of which $\mathcal{B}$ is a basin. Therefore, the cycle is contained in $\partial\Delta$ and cannot be attracting so it is parabolic. Unfortunately the separation theorem does not prevent this: the separating external rays could as well land on this parabolic point and still separate $\mathcal{B}$ from Δ.

This is the only place in the proof of case (3) where we use the assumption $n_c = 2$.

4.7.8. *Hyperbolicity.* Let us cite Theorem A in [Mañ85] (see also [Mañ87]).

Theorem (Mañé)**.** *Let $N = S^1$ (the circle) or $N = [0, 1]$. If f is a C^2 map from N to N and $\Lambda \subset N$ is a compact invariant set that doesn't contain critical points, sinks or non-hyperbolic periodic points, then either $\Lambda = N = S^1$ and f is topologically equivalent to a rotation or Λ is a hyperbolic set.*

By the preceding discussion, we can apply this to $\widetilde{f}$ on $\partial\mathbb{D}$ with $N = \partial\mathbb{D}$: Section 4.7.7 proved the absence of attracting or parabolic cycle, and Lemma 18 the absence of critical points.

So $\widetilde{f}$ is hyperbolic on $\partial\mathbb{D}$, which means that there is a continuous function $\rho(z) > 0$ on the unit circle such that $\widetilde{f}$ is uniformly expanding with respect to the metric $\rho(z)|dz|$, i.e. $\rho(\widetilde{f}(z))|\widetilde{f}'(z)| > \kappa\rho(z)$ for some $\kappa > 1$. Consider for some $\varepsilon > 0$ the domain

$$V = \bigcup_{\theta\in\mathbb{R}} \left(\exp(i\theta - \varepsilon\rho(e^{i\theta})), \exp(i\theta + \varepsilon\rho(e^{i\theta}))\right).$$

Let U be the connected component U of $\widetilde{f}^{-1}(V)$ containing the unit circle. For ε small enough, U is compactly contained in V.

Let $U' = \widehat{\Delta}\cup\phi^{-1}(U)$ and $V' = \widehat{\Delta}\cup\phi^{-1}(V)$. Then U' and V' are open, connected, simply connected, U' is a connected component of $P^{-p}(V')$ and U' is compactly contained in V'. Therefore the restriction of $P^p : U' \to V'$ is a polynomial like map. This restriction is also unicritical.

So we have proved that P^p has a unicritical polynomial like restriction whose domain contains Δ. For the restriction, Δ is a Siegel disk of rotation number Δ and period one. Now we can either apply Herman's theorem (which works as well for unicritical polynomial like maps): the critical point c' must belong to $\partial\Delta$. This contradict our standing assumption that $P(c') \notin \partial\Delta$. Or instead we can conclude as in case 2: Fatou's theorem stating that $\partial\Delta$ is contained in the ω-limit set of the critical points also works for polynomial-like maps, and since the only critical point c' of our polynomial like map is in a Fatou component, it cannot accumulate on all of $\partial\Delta$, leading to a contradiction.

This finishes the proof of the second case: the assumption $P(c') \notin \partial\Delta$ leads to a contradiction.

References

[Ché11] Arnaud Chéritat. Relatively compact Siegel disks with non-locally connected boundaries. *Math. Ann.*, 349(3):529–542, 2011.

[GŚ03] Jacek Graczyk and Grzegorz Świątek. Siegel disks with critical points in their boundaries. *Duke Math. J.*, 119(1):189–196, 2003.

[Her79] Michael-Robert Herman. Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Inst. Hautes Études Sci. Publ. Math.*, (49):5–233, 1979.

[Her85] Michael-R. Herman. Are there critical points on the boundaries of singular domains? *Comm. Math. Phys.*, 99(4):593–612, 1985.

[Kiw00] Jan Kiwi. Non-accessible critical points of Cremer polynomials. *Ergodic Theory Dynam. Systems*, 20(5):1391–1403, 2000.

[Mañ85] Ricardo Mañé. Hyperbolicity, sinks and measure in one-dimensional dynamics. *Comm. Math. Phys.*, 100(4):495–524, 1985.

[Mañ87] Ricardo Mañé. Erratum: "Hyperbolicity, sinks and measure in one-dimensional dynamics". *Comm. Math. Phys.*, 112(4):721–724, 1987.

[Mañ93] Ricardo Mañé. On a theorem of Fatou. *Bol. Soc. Brasil. Mat. (N.S.)*, 24(1):1–11, 1993.

[Mil06] John Milnor. *Dynamics in one complex variable*, volume 160 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, third edition, 2006.

[Rog98] James T. Rogers, Jr. Diophantine conditions imply critical points on the boundaries of Siegel disks of polynomials. *Comm. Math. Phys.*, 195(1):175–193, 1998.

[Yoc02] Jean-Christophe Yoccoz. Analytic linearization of circle diffeomorphisms. In *Dynamical systems and small divisors (Cetraro, 1998)*, volume 1784 of *Lecture Notes in Math.*, pages 125–173. Springer, Berlin, 2002.

[Zha08] Gaofei Zhang. All bounded type Siegel disks of rational maps are quasi-disks, 2008. arXiv:0810.2733.

Centre National de la Recherche Scientifique, Institut de Mathématiques de Toulouse, Université Paul Sabatier 118 Route de Narbonne 31062 Toulouse Cedex 9, France
E-mail address: `arnaud.cheritat@math.univ-toulouse.fr`

Institut de Mathématiques de Toulouse, Université Paul Sabatier 118 Route de Narbonne 31062 Toulouse Cedex 9, France
E-mail address: `pascale.roesch@math.univ-toulouse.fr`